

# eScholarship@UMassChan

## Impact of medical and neurological ICU complications on moderate-severe traumatic brain injury (TBI)

Item Type	Poster Abstract
Authors	Muehlschlegel, Susanne;Carandang, Raphael A.;Ouillette, Cynthia;Hall, Wiley R.;Anderson, Frederick A. Jr.;Goldberg, Robert J.
DOI	<a href="https://doi.org/10.13028/3td9-m949">10.13028/3td9-m949</a>
Rights	Copyright the Author(s)
Download date	2024-12-31 10:26:24
Item License	<a href="http://creativecommons.org/licenses/by-nc-sa/3.0/">http://creativecommons.org/licenses/by-nc-sa/3.0/</a>
Link to Item	<a href="https://hdl.handle.net/20.500.14038/27756">https://hdl.handle.net/20.500.14038/27756</a>

## **IMPACT OF MEDICAL AND NEUROLOGICAL ICU COMPLICATIONS ON MODERATE-SEVERE TRAUMATIC BRAIN INJURY (TBI)**

Susanne Muehlschlegel, MD, MPH<sup>1,2,3</sup>; Raphael Carandang, MD<sup>1,3</sup>; Cynthia Ouillette, RN<sup>1</sup>; Wiley Hall, MD<sup>1,3</sup>; Fred Anderson, PhD<sup>3,4</sup>; Robert Goldberg, PhD<sup>5</sup>

Departments of <sup>1</sup>Neurology (Div. Neurocritical Care), <sup>2</sup>Anesthesia/Critical Care and <sup>3</sup>Surgery; <sup>4</sup>Center for Outcomes Research; <sup>5</sup>Department of Quantitative Health Sciences (Div. Epidemiology of Chronic Diseases), University of Massachusetts Medical School, Worcester, MA

### Contact information:

Susanne Muehlschlegel, MD, MPH

Email: susanne.muehlschlegel@umassmemorial.org

Phone: 508-856-4684

### **Abstract:**

Certain admission characteristics are known predictors of adverse outcomes in patients with moderate-severe TBI, but explain only 1/3 of outcome variability. Intensive care unit (ICU) complications occur frequently in this population, but their impact on patient outcomes remains poorly defined. In a prospective observational cohort study of 170 consecutive moderate-severe TBI patients admitted to Level I trauma center (UMASS) over the period 11/2009–2/2012, we examined the association of ICU complications and 3-month outcome (Glasgow Outcome Scale [GOS]). The mean age was 51 years, 72% were men, and the median GCS and injury severity scores were 4 and 29, respectively. Using multiple logistic regression analysis, hypotension requiring vasopressors (HRV) was the strongest predictor of poor outcome (GOS 1-3 [OR 2.8; 95% CI 1-7.5]) among medical complications. After combining medical with neurological ICU complications, brain herniation (OR 5.8; 95% CI 1.1-30.2) and intracranial rebleeding (OR 2.9; 95% CI 1-8.4) were the strongest predictors of poor outcome, while HRV approached significance (OR 2.4; 95% CI 0.9-6.4). We identified important potentially modifiable predictors of adverse outcomes after moderate-severe TBI. Confirmation of our findings in a larger cohort is warranted.