

Nuclear Export Signal and Immunodominant CD8⁺ T Cell Epitope in Influenza A Virus Matrix Protein 1

Shuai Cao et al. recently identified and characterized a nuclear export signal (NES) in influenza A virus matrix protein 1 (M1) (5). We noticed that the identified NES, ⁵⁹ILGFVFTLTV⁶⁸, almost completely overlaps the HLA-A2-restricted CD8⁺ T cell epitope, ⁵⁸GILGFVFTL⁶⁶ (6), which is considered immunodominant in individuals carrying the HLA-A2 allele (3, 15).

Colocalization of the NES and the immunodominant HLA-A2-restricted CD8⁺ T cell epitope in M1 explains why escape mutation in this epitope was not found in nature. Escape mutation from CD8⁺ T cell recognition are known for the nucleoprotein of influenza A virus in the context of HLA-B*08, -B*2705, and -B*3501 alleles (4, 11, 13). Considering that the HLA-A*0201 allele or alleles belonging to the HLA-A2 supertype are very common in the human population (more than 40%) (12), it is expected that escape mutation may emerge in the M1 CD8⁺ T cell epitope. However, as Cao et al. pointed out in their paper, the NES in M1 is highly conserved among influenza A subtypes (5). Berkhoff et al. previously showed that mutation at the P2 anchor residue in the M1 epitope (the I residue at position 59) costs viral fitness (1, 2). Now we know that the fitness cost is due to the loss of functional NES in M1.

Another implication of the colocalization is that there may be more cases of HLA-A2-restricted CD8⁺ T cell epitopes overlapping with the NES in viral or cellular proteins. Key residues in the NES (ILGFVFTLTV) are also anchor or auxiliary anchor residues (in bold type) in the HLA-A2 binding motif (GILGFVFTL) (10) except for the V at position 68, which is outside of the CD8⁺ T cell epitope. We searched for the HLA-A*0201 binding motif overlapping with the NES mentioned in the paper by Cao et al. (in Table 1 and the text of the article [5]) using the epitope prediction algorithms, HLA Peptide Binding Predictions at BIMAS (http://www-bimas.cit.nih.gov/molbio/hla_bind/) (8) and SYFPEITHI (<http://www.syfpeithi.de/>) (9). Among them two NESs were found to overlap with the HLA-A*0201 binding motifs. The NES in the receptor-interacting protein 3 (RIP3), ¹¹⁶LLCRLKKEVVL¹²⁶ (16) with ¹¹⁶LLCRLKKEV¹²⁴ (the BIMAS score is 271.948 and the SYFPEITHI score is 28, while the scores for ⁵⁸GILGFVFTL⁶⁶ are 550.927 and 30, respectively), and the NES of Nipah virus matrix protein, ¹⁰⁶LLEELCSLKV¹¹⁵ (14), with ¹⁰⁵DLEELCSL¹¹³ (the BIMAS score is 55.902, and the SYFPEITHI score is 29). Colocalization of the NES and the HLA-A*0201-restricted epitopes is consistent with recent bioinformatic analysis by Hertz et al. showing that human major histocompatibility complex (MHC) class I molecules tend to target conserved regions of human and viral proteins (7).

REFERENCES

- Berkhoff EG, et al. 2006. Fitness costs limit escape from cytotoxic T lymphocytes by influenza A viruses. *Vaccine* 24:6594–6596.
- Berkhoff EG, et al. 2005. Functional constraints of influenza A virus epitopes limit escape from cytotoxic T lymphocytes. *J. Virol.* 79:11239–11246.
- Boon AC, et al. 2002. The magnitude and specificity of influenza A virus-specific cytotoxic T-lymphocyte responses in humans is related to HLA-A and -B phenotype. *J. Virol.* 76:582–590.
- Boon AC, et al. 2002. Sequence variation in a newly identified HLA-B35-restricted epitope in the influenza A virus nucleoprotein associated with escape from cytotoxic T lymphocytes. *J. Virol.* 76:2567–2572.
- Cao S, et al. 2012. A nuclear export signal in the matrix protein of influenza A virus is required for efficient virus replication. *J. Virol.* 86:4883–4891.
- Gotch F, McMichael A, Rothbard J. 1988. Recognition of influenza A matrix protein by HLA-A2-restricted cytotoxic T lymphocytes. Use of analogues to orientate the matrix peptide in the HLA-A2 binding site. *J. Exp. Med.* 168:2045–2057.
- Hertz T, et al. 2011. Mapping the landscape of host-pathogen coevolution: HLA class I binding and its relationship with evolutionary conservation in human and viral proteins. *J. Virol.* 85:1310–1321.
- Parker KC, Bednarek MA, Coligan JE. 1994. Scheme for ranking potential HLA-A2 binding peptides based on independent binding of individual peptide side-chains. *J. Immunol.* 152:163–175.
- Rammensee H, Bachmann J, Emmerich NP, Bachor OA, Stevanovic S. 1999. SYFPEITHI: database for MHC ligands and peptide motifs. *Immunogenetics* 50:213–219.
- Rammensee HG, Friede T, Stevanovic S. 1995. MHC ligands and peptide motifs: first listing. *Immunogenetics* 41:178–228.
- Rimmelzwaan GF, et al. 2004. Sequence variation in the influenza A virus nucleoprotein associated with escape from cytotoxic T lymphocytes. *Virus Res.* 103:97–100.
- Sette A, Sidney J. 1999. Nine major HLA class I supertypes account for the vast preponderance of HLA-A and -B polymorphism. *Immunogenetics* 50:201–212.
- Voeten JT, et al. 2000. Antigenic drift in the influenza A virus (H3N2) nucleoprotein and escape from recognition by cytotoxic T lymphocytes. *J. Virol.* 74:6800–6807.
- Wang YE, et al. 2010. Ubiquitin-regulated nuclear-cytoplasmic trafficking of the Nipah virus matrix protein is important for viral budding. *PLoS Pathog.* 6:e1001186. doi:10.1371/journal.ppat.1001186.
- Wu C, et al. 2011. Systematic identification of immunodominant CD8⁺ T-cell responses to influenza A virus in HLA-A2 individuals. *Proc. Natl. Acad. Sci. U. S. A.* 108:9178–9183.
- Yang Y, Ma J, Chen Y, Wu M. 2004. Nucleocytoplasmic shuttling of receptor-interacting protein 3 (RIP3): identification of novel nuclear export and import signals in RIP3. *J. Biol. Chem.* 279:38820–38829.

Masanori Terajima

Francis A. Ennis

Center for Infectious Disease and Vaccine Research
Division of Infectious Diseases and Immunology
Department of Medicine
University of Massachusetts Medical School
Worcester, Massachusetts, USA

Address correspondence to Masanori Terajima,
Masanori.Terajima@umassmed.edu.

For the author reply, see doi:10.1128/JVI.01245-12.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.
doi:10.1128/JVI.00894-12