

The *C. elegans* Snail homolog CES-1 can activate gene expression *in vivo* and share targets with bHLH transcription factors

John S. Reece-Hoyes¹, Bart Deplancke¹, M. Inmaculada Barrasa¹, Julia Hatzold², Ryan B. Smit³, H. Efsun Arda¹, Patricia A. Pope¹, Jeb Gaudet³, Barbara Conradt² and Albertha J. M. Walhout^{1,*}

¹Program in Gene Function and Expression and Program in Molecular Medicine, University of Massachusetts Medical School, Worcester, MA 01605, USA, ²Department of Genetics, Dartmouth Medical School, Hanover, NH 03755, USA and ³Department of Biochemistry & Molecular Biology, University of Calgary, Health Sciences Centre, Calgary, Alberta T2N 4N1, Canada

Received February 9, 2009; Revised March 25, 2009; Accepted March 28, 2009

ABSTRACT

Snail-type transcription factors (TFs) are found in numerous metazoan organisms and function in a plethora of cellular and developmental processes including mesoderm and neuronal development, apoptosis and cancer. So far, Snail-type TFs are exclusively known as transcriptional repressors. They repress gene expression by recruiting transcriptional co-repressors and/or by preventing DNA binding of activators from the basic helix-loop-helix (bHLH) family of TFs to CAGGTG E-box sequences. Here we report that the *Caenorhabditis elegans* Snail-type TF CES-1 can activate transcription *in vivo*. Moreover, we provide results that suggest that CES-1 can share its binding site with bHLH TFs, in different tissues, rather than only occluding bHLH DNA binding. Together, our data indicate that there are at least two types of CES-1 target genes and, therefore, that the molecular function of Snail-type TFs is more plastic than previously appreciated.

INTRODUCTION

Snail-type TFs have been identified in numerous metazoan organisms, first in the fruit fly *Drosophila melanogaster* (1), and subsequently in others, including the nematode *Caenorhabditis elegans* (2), and human (3,4).

Snail-type TFs are involved in many biological processes such as development, mesodermal and neuronal differentiation, apoptosis and cancer (5–7). The Snail-type family of TFs includes members that most resemble the firstly identified Snail, and proteins that resemble the homologous proteins Scratch and Slug. They possess three to five C2H2 zinc fingers with which they bind DNA in a sequence-specific manner. The DNA-binding specificity of several Snail-type TFs has been characterized to a limited extent. *In vitro* gel shift assays (2,8–11) and recently, *in vivo* (12) studies found that the Snail-type TFs bind the CAGGTG E-box and often more specifically the ACAGGTG sequence (referred to here as ‘Snail-box’).

So far, Snail-type TFs have been exclusively reported to function as repressors *in vivo* [see e.g. (3,4,13–15)]. The repressor activity of vertebrate Snail-type TFs depends on the SNAG repressor domain, which mediates repression by recruiting histone deacetylases (6,16). However, except for Scratch, *C. elegans* and *Drosophila* Snail-type TFs do not contain a SNAG domain. Other mechanisms of repression by Snail-type TFs include interactions with the co-repressors dCtBP (17) and Ebi (18). In addition, Snail-like TFs can repress gene expression by preventing the binding of members of the basic region helix-loop-helix (bHLH) family of TFs to DNA, thereby antagonizing the function of these transcriptional activators (3,10,13,15,19).

The *C. elegans* genome encodes three Snail-type C2H2 TFs: CES-1, K02D7.2 and SCRT-1 (6) (Figure 1a). A gain-of-function (gf) in *ces-1* (cell death specification-1)

*To whom correspondence should be addressed. Tel: +1 508 856 4364; Fax: +1 508 856 5460; Email: marian.walhout@umassmed.edu
Present address:

Bart Deplancke, Institute of Bio-engineering, Ecole Polytechnique Fédérale de Lausanne, Lausanne, CH-1015, Switzerland.

The authors wish it to be known that, in their opinion, the first two authors should be regarded as joint First Authors.

© 2009 The Author(s)

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/2.0/uk/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

was first discovered in a genetic screen to identify genes involved in programmed cell death (20). The *ces-1* gene product was found to be most homologous to the Scratch members of the Snail superfamily (2). Like other Snail-type TFs, CES-1 has been reported to function as a transcriptional repressor. Specifically, in *ces-1(gf)* animals, CES-1 is overexpressed and represses its only known target gene, *egl-1*, by preventing binding of activating HLH-2/HLH-3 heterodimers to the multiple E/Snail-boxes that are located in or around this target gene (15). So far no transcriptional effect has been reported in *ces-1* loss-of-function (*lf*) animals.

Surprisingly, we find that CES-1 can function as a transcriptional activator *in vivo*, by binding to a single Snail-box in the promoter of *B0507.1*. We find that CES-1 binds an extended Snail-box that is comprised of nine nucleotides and which we refer to as a 'CES-box'. Finally, our findings strongly suggest that CES-1 shares rather than antagonizes the target *B0507.1* promoter with HLH-6, another bHLH family member, although each TF utilizes the site in different tissues. Thus, there are at least two types of CES-1 target genes: those that are repressed by CES-1 through a bHLH occlusion mechanism and those that are activated by both types of TFs, but in different tissues. Taken together, the transcriptional function of Snail-type TFs as well as their functional interactions with other TFs may be more flexible than previously thought.

MATERIALS AND METHODS

Yeast one-hybrid assays

Yeast one-hybrid (Y1H) assays were performed as described previously (21–23).

Mutant promoter and binding site constructs

Mutant promoter constructs were generated by PCR using plasmid templates and primers listed in Table S1. Mutant *Promoter::GFP* constructs were generated by mutating the promoter Entry vector construct and transferring the changed insert to pDD04 (24) by a Gateway LR recombination reaction (25). pENTRY::*PB0507.1*(Δ CES-box) was created using overlapping primers and the QuickChange Site Directed Mutagenesis Kit (Stratagene) that generates a mutant circular plasmid. All other mutant constructs were created using non-overlapping primers and PCR amplification with Platinum Pfx (Invitrogen) generating linear PCR products that require excision and extraction from agarose gel (Qiagen kit), treatment with PNK (New England Biolabs) and ligation using T4 Ligase (New England Biolabs). Binding site constructs were generated by cloning a DNA fragment containing a single copy of the putative binding site into BamHI/HindIII-digested Gateway-compatible Entry vector pMW#4 that we created for binding site cloning. These DNA fragments were created by annealing complementary primers (Table S1) with restriction enzyme-compatible overhangs. All constructs were confirmed by sequencing (Agencourt).

Caenorhabditis elegans transgenesis and mutant crosses

Transgenic animals were created by ballistic transformation in *unc-119(ed3)* animals as described (26,27). For each GFP construct we obtained multiple (up to eight) independent lines that all exhibited identical GFP expression patterns. Frozen stocks were generated for a maximum of three of these lines, and the best transmitting line was used for all experiments. For *PB0507.1::GFP* the strain VL456 was used because it contains the transgene integrated in the genome. For *PB0507.1*(Δ CES-box)::GFP and *PB0507.1*(A+1C)::GFP, the non-integrated strains VL457 and VL708 were used respectively. Both VL457 and VL708 exhibited a >90% transmittal rate. The loss-of-function mutant strains *ces-1* (*n703;n1434*) (2), *K02D7.2(bc366)*, *scrt-1(tm509)* and *hlh-6(tm299)* (28) were crossed into VL456.

Caenorhabditis elegans imaging and GFP expression scoring

Immobilized worms were viewed using a Zeiss Axioscope 2 and images were acquired using a Hamamatsu Orca-ER Digital Camera. GFP expression was scored for each genotype in least 100 worms (50 L1 to L3 larval stages, and 50 L4 to adult).

Bioinformatics

The sequence logo for the CES-1 binding site was created using <http://weblogo.berkeley.edu/> (29). To search for promoters containing the CES-1-binding site we created a position weight matrix (PWM) using sequences from –5 to +7 (Figure 2a) from the five promoters initially found to bind CES-1 in our Y1H assays. We searched all *C. elegans* promoter sequences (24) (WS93) using Improbizer motif matcher (30) with this PWM. We considered promoters with an Improbizer score of 10.4 or higher as having a putative CES-1-binding site; this cut-off was defined by looking at the scores of the five Y1H positive promoters. For Y1H assays we selected 24 promoters for which a clone was available in the Promoterome (24).

RESULTS

CES-1 binds a single Snail-type DNA element

Interactions between TFs and their target genes can be modeled into transcription regulatory networks that provide insights into gene expression at a systems level (31). We have previously used Y1H assays to map transcription regulatory networks that pertain to the *C. elegans* digestive tract, nervous system and microRNA regulation (32–34). In each of these networks, we retrieved the Snail-type TF CES-1 with one or more promoters. These promoters include *PB0507.1*, *Plin-32*, *Pmir-235* and *Pmir-231*. We recently also mapped a transcription regulatory network pertaining to fat biology and, in that study, have retrieved CES-1 with the promoter of *C30F12.1* (H.E.A. and A.J.M.W., in preparation). We obtained clones encoding CES-1 and its homologs K02D7.2 and SCRT-1 from our TF array resource (23) to examine the




Figure 1. A single site is necessary and sufficient for CES-1 DNA binding. (a) The *C. elegans* Snail family contains three members: CES-1, K02D7.2 and SCRT-1. The C2H2 zinc fingers are indicated by purple boxes. Number in each of the boxes indicates percentage identity to the corresponding CES-1 zinc finger. (b) CES-1 binds *PB0507.1*, *Pmir-235*, *Plin-32*, *PC30F12.1* and *Pmir-231* in Y1H assays, whereas its closest homologs, K02D7.2 and SCRT-1 do not. P—permissive media; H—selective media lacking histidine, 20mM 3-aminotriazole; Z—colorimetric assay for *LacZ* activity; AD—Gal4p activation domain. (c) Each of the five promoters that bind CES-1 possesses a Snail-box or Snail-box-like motif. Promoter length is indicated at the left; position, strand (+) or (-), and sequence are indicated at the right. Red—conventional Snail-box sequence (ACAGGTG), blue—Snail-box-like sequence (ACAGGTT). (d) Y1H assays showing that CES-1 cannot bind promoters in which the Snail-box or Snail-box-like motif is deleted. ZTF-2 binding to *PB0507.1* and TBX-8 binding to *Plin-32* are shown as positive controls for the functionality of mutant promoter constructs. The mutant yeast strains contain a wild-type promoter::*LacZ* construct to ensure functionality of AD-CES-1. (e) Y1H assays showing that a single copy of a CES-box is sufficient for CES-1 binding. Both the *PB0507.1* and *Plin-32* CES-box variants are shown.

specificity of CES-1 DNA binding. We found that neither K02D7.2 nor SCRT-1 (Figure 1b) was capable of binding these promoters in Y1H assays, indicating that the interaction between CES-1 and the five promoters is specific.

Like its evolutionary counterparts, CES-1 has been reported to interact with the E/Snail-box, (A)CAGGTG (2). We examined the sequence of each of the promoters that interact with CES-1 and found that four of these indeed contain a Snail-box (Figure 1c). *Plin-32* does not

possess a perfect Snail-box but does contain a sequence that differs from a Snail-box only in a single nucleotide (Snail-box-like sequence, ACAGGTT) (Figure 1c).

To test whether the Snail-box and Snail-box-like sequences function as CES-1-binding sites we deleted them from each promoter and examined CES-1 binding by Y1H assays. We deleted the putative CES-1-binding sites only in the context of the *Promoter::HIS3* constructs and used wild-type *Promoter::LacZ* constructs. This enabled us to verify the functionality of CES-1 in yeast. As expected, we found that CES-1 failed to bind promoters in which the Snail-box was deleted, demonstrating that this sequence is required for CES-1 binding (Figure 1d, left panel, Figure S1 and data not shown). When we deleted the CAGGTT Snail-box-like sequence from *Plin-32*, we found that CES-1 binding is similarly abolished (Figure 1d, right panel, see also below). Importantly, interactions with other TFs were not affected upon the deletion of the putative CES-1-binding sites, which demonstrates that each mutated promoter was functional in Y1H assays (Figure 1d and Figure S1).

Each of the CES-1-interacting promoters contains a single putative CES-1-binding site, suggesting that one site may be sufficient for CES-1 binding. To test this, we cloned each of the two different sites into our Y1H vectors and generated Y1H bait strains (21,22). We found that CES-1 is indeed capable of binding a single site, and can bind both Snail-box variants (Figure 1e). This demonstrates that a single site is sufficient for CES-1 binding. Additionally, the CES-binding site can occur on either strand (Figure 1c), which indicates that CES-1 binding is independent of binding site orientation.

CES-1 binds an extended Snail-box

Our finding that CES-1 can bind two different sites, CAGGTG and CAGGTT, suggests that CES-1 may have a broader DNA-binding specificity than previously reported. However, specificity of binding may also be influenced by sequences outside the core Snail-box, as CES-1 contains five C2H2 zinc fingers, each of which may contact up to three nucleotides upon binding of the protein to DNA (Figure 1a) (35). We used the CES-1-binding site sequences and flanking nucleotides from each of the five promoters that were bound by CES-1 to derive a CES-1-binding-site logo (Figure 2a). We identified two additional positions that were identical or highly similar in these promoters, all flanking the 5' end of the Snail-box or Snail-like-box. The -3 position was a C in all cases and the -2 position was either a C or an A. We refer to this longer CES-1-binding site as the 'CES-box' (Figure 2a).

To test the contribution of each nucleotide to CES-1 binding, we created mutant CES-box constructs, all in the context of the *B0507.1* promoter. We verified the integrity and functionality of each construct by sequencing, and by ensuring that the interaction with a non-related TF (ZTF-2) was unaffected (Figure 2b). We found that all positions in the CES-box are important for CES-1 binding, including positions -3 and -2. Surprisingly, changing the G in position +6 to a T also

abolished CES-1 binding in the context of *PB0507.1*, even though that site now exactly matched that found in *Plin-32* (Figure 2b). This suggests that promoter context may influence either the specificity or the affinity of CES-1 DNA binding. For example, nucleotides at the -4, +7 and +8 positions could influence the affinity of CES-1 for the *Plin-32* site. Taken together, our DNA-binding specificity analysis revealed an extended CES-1-binding site that is composed of nine nucleotides and that contains a Snail box or Snail-box-like sequence.

Predictive power of the CES-1-binding site

To determine the predictive power of the different CES-1-binding sites, we searched all predicted *C. elegans* gene promoters (24) for the previously reported Snail-binding sites: the CAGGTG E-box and the ACAGGTG Snail-box, as well as the newly defined CES-box that is longer and that can tolerate a T at the +6 position (Figure 2a). Out of 20441 promoters examined, we found the CAGGTG E-box in 5250 (26%), the Snail-box in 1896 (9%) and the CES-box in 406 (2%) promoters (Table 1). Next, we examined the promoters that we had previously tested for binding to CES-1 in Y1H assays (32–34; H.E.A. and A.J.M.W., in preparation). We found that, of the Y1H positives that interact with CES-1, 80% contained an E-box, 80% contained a Snail-box and 100% contained a CES box sites, compared to 35%, 11% and 2% of the Y1H negatives (Table 1). Conversely, we found that 4 out of 87 (5%) of the promoters harboring a CAGGTG E-box tested interacted with CES-1, compared to 4 out of 29 (14%) of the promoters with a Snail-box and 5 out of 10 (50%) of the promoters with a CES-box (Table 1). Using a Fisher exact test we determined that the association between the presence of a site and the binding of CES-1 in Y1H assays is more significant for the CES-box ($P < 10^{-7}$) than either the E-box ($P = 0.057$) or Snail-box ($P < 10^{-3}$) (Table 1). This demonstrates that the CES-box is a better predictor of CES-1 binding than either of the shorter sequences. As a control we used the reverse (but not complement) sequences that have similar nucleotide composition, and did not observe an increase in observed over expected occurrence in promoters that bind CES-1 (Table 1).

We have previously created a *C. elegans* 'Promoterome' resource that consists of ~6000 Gateway-cloned protein-coding gene promoters, complemented by ~75 microRNA promoters (24,36). We identified which of the cloned promoters contain CES-box, and selected 24 to test for CES-1 binding in Y1H assays. Of these 24, eight possess CAACAGGTG, five have CCACAGGTG, six have CAACAGGT T and five have CCACAGGTT (Figure 2c). We created Y1H DNA bait strains for all 24 promoters and transformed in AD-CES-1 plasmid and an AD-alone plasmid (negative control). Surprisingly, only six of these promoters were capable of interacting with CES-1 (Figure 2c, Figure S2). This suggests that additional nucleotides surrounding the extended CES-box may influence CES-1 binding. However, by examining the nucleotides flanking the 5' and 3' end of the CES-box, we could not find




Figure 2. CES-1 binds an extended Snail-box. (a) Logo of the CES-box, an extended Snail-box, based on CES-1/promoter interactions identified by Y1H assays. (b) Y1H assays examining CES-1 binding to *PB0507.1* variants that carry point mutations in the CES-box. AD—negative control with only the Gal4p activation domain; AD-ZTF-2—positive control demonstrating the functionality of the mutant baits. (c) Six out of 24 promoters that contain a CES-box interact with CES-1 in Y1H assays (indicated in red).

obvious candidates. Alternatively, not all CES-boxes may be accessible due to nucleosome positioning. The latter may be more likely since Y1H baits are integrated into the yeast genome at a fixed location and copy number, and are therefore embedded into chromatin (21). Interestingly, CES-1 bound to 50% of the tested promoters with a *PB0507.1*-like CAACAGGTG motif. However, CES-1 bound only one of five promoters with a *Plin-32*-like CAACAGGTT motif, one of six with a *Pmir-235*-like CCACAGGTG motif, and none of five with the CCACAGGTT sequence that we did not observe in any of our original Y1H positive promoters (Figure 1c, Figure S2). This suggests that the latter may not constitute

a functional CES-1-binding site and reflects the loss of information in binding site logos that compile multiple data points into a single model, but that do not reflect combinations of nucleotides that are avoided. Although the differences observed between the sites are based on small samples, these results indicate that the various CES-box variants may provide different degrees of predictive power.

CES-1 activates promoter activity *in vivo*

To examine the effect of CES-1 on *C. elegans* promoter activity *in vivo*, we focused on the 300 bp promoter of

Table 1. Occurrence of CES-1-binding sites

	E-box	Snail-box	CES-box	Reverse E-box	Reverse Snail-box	Reverse CES-box
Promoters with hits (20 441)	5250 (26%)	1896 (9%)	406 (2%)	3543 (17%)	1420 (7%)	457 (2%)
Y1H neg promoters (238)	83 (35%)	25 (11%)	5 (2%)	53 (22%)	27 (11%)	6 (3%)
Y1H pos promoters (5)	4 (80%)	4 (80%)	5 (100%)	1 (20%)	0	0
P-value (Fisher exact test)	0.057	<10 ⁻³	<10 ⁻⁷	1	0.624	1




Figure 3. CES-1 and its binding site activate *PB0507.1* in vivo. (a) Images of GFP expression in *PB0507.1::GFP* animals. The 300 bp promoter of *B0507.1* drives GFP expression in the pharyngeal glands (PG), pharyngeal-intestinal valve (PIV), spermatheca (SPT), rectal gland (RG) and distal tip cell (DTC) (indicated by arrow heads). Left—GFP expression; center—DIC image; right—merge. (b) Loss of *K02D7.2* has no effect on *PB0507.1* activity. (c) Deletion or point mutations in the CES-box in *PB0507.1* result in different changes in expression in the PG. Top—GFP expression; center—DIC image; bottom—merge. (d) Quantification of GFP expression data in TF or TF binding site mutants. PPGC—posterior pharyngeal gland cells; APGC—anterior pharyngeal gland cells.

B0507.1. We used Gateway cloning (25,38) to create a transcriptional fusion with *PB0507.1* upstream of an open reading frame (ORF) encoding the green fluorescent protein (GFP), and used the resulting construct to create transgenic *C. elegans* that carry the transgene integrated into the genome (27,36). Subsequently, we analyzed spatiotemporal promoter activity by examining where and when GFP is expressed in these animals. We found that *PB0507.1* drives GFP expression in all developmental stages and in multiple tissues, including the anterior and posterior pharyngeal gland cells, pharyngeal-intestinal valve, rectal gland, spermatheca and distal tip

cells (Figure 3a). This expression pattern was identical in multiple independent transgenic lines (data not shown). To test whether CES-1 and/or its closest homologs regulate the activity of *PB0507.1* in vivo, we crossed the *PB0507.1::GFP* animals with animals carrying a deletion in *ces-1*, *K02D7.2* or *srt-1*. Surprisingly, we found that part of the expression pattern disappears upon removal of CES-1; while expression was maintained in the pharyngeal glands, expression in the pharyngeal-intestinal valve, spermatheca, distal tip cell and rectal gland was greatly reduced (Figure 3a, quantification in Figure 3d). This suggests that CES-1 functions as an activator of *B0507.1*

promoter activity. The activity of *PB0507.1* was unchanged in *K02D7.2(bc366)* (Figure 3b and d) and *scrt-1(ok1228)* mutants (data not shown), which is in agreement with the observation that these CES-1 homologs cannot interact with this promoter in Y1H assays (Figure 1b).

We examined whether we could detect an overlap in *Pces-1* and *PB0507.1* activity but did not observe a clear overlap in later stages of development (38; data not shown). However, the promoter fragments used may be missing the regulatory elements required for such expression, or do not drive robust enough expression for the overlap to be observed. We did observe broad *ces-1* promoter activity in embryos (38; data not shown). However, it is difficult to assess whether this activity overlaps with that of *PB0507.1*. Thus, it is formally possible that CES-1 activates the *B0507.1* promoter indirectly, for instance by repressing a transcriptional repressor. To test this, we examined the function of the *PB0507.1* CES-box *in vivo*. We fused *B0507.1* promoter variants that carry a CES-box deletion or point mutation (Figures 1d and 2b) to the GFP-encoding ORF by Gateway cloning and created transgenic worms. Interestingly, we found different GFP expression patterns with the CES-box deletion construct compared to the CES-box point mutant (-1A to C) (Figure 1c), neither of which can interact with CES-1 (Figures 1d and 2b). When the CES-box was completely removed, GFP expression disappeared in all tissues/cells, but remained unchanged in the posterior pharyngeal gland cells (Figure 3c and d). With the CES-box point mutant, however, we only observed a partial reduction of GFP expression in the anterior pharyngeal glands and spermatheca, and lack of expression in the pharyngeal-intestinal valve and rectal gland (Figure 3c and d). The expression in the posterior pharyngeal glands was unaffected in the substitution mutant, similar to that observed in the CES-box deletion mutant. Together these observations indicate that CES-1 activates *PB0507.1* by binding the CES-box. Additionally, the observation that the CES-box deletion mutant has a broader and stronger effect on *PB0507.1* activity, and that expression of GFP in the anterior pharyngeal glands is (partially) reduced in the CES-box mutants, but not in *ces-1(lf)* animals indicates that at least one additional factor activates *PB0507.1* through the CES-box. Finally, the fact that expression in the posterior pharyngeal glands is unaffected in either *ces-1(lf)* or CES-box mutants suggests that another regulator independently activates *PB0507.1* in this tissue by binding to another *cis*-regulator element in the promoter.

HLH-6 activates *PB0507.1* in the anterior pharyngeal glands

Snail-type TFs have been reported to antagonize transcriptional activation by bHLH TFs by binding to the same *cis*-regulatory DNA elements (3,13,15,19). bHLH TFs are known to bind CANNTG E-box sequences. The promoter of *hlh-6* specifically drives expression in the pharyngeal glands (27,28,39), and the HLH-6 protein is involved in transcriptional regulation in this tissue (28).

Therefore, we hypothesized that HLH-6 may activate *PB0507.1* in the pharyngeal glands. We crossed *PB0507.1::GFP* worms with animals that carry a deletion in the *hlh-6* gene [*hlh-6(tm299)*] (28), and found that GFP expression was reduced in the anterior pharyngeal glands, but unaffected in one of the three posterior pharyngeal glands (g1P) in a subset of the animals. The other two posterior glands (g2R and g2L) are absent in ~80% of *hlh-6(tm299)* mutants (Figure 4a and b). HLH-6 specifically dimerizes with HLH-2, the ortholog of E12/E47 and Daughterless (C. Grove and A.J.M.W., in preparation). However, since *hlh-2* loss-of-function leads to lethality (40), we could not examine the effect of HLH-2 on *PB0507.1* activity. Taken together, our data suggest that HLH-6 may share its DNA target with CES-1 in different tissues rather than antagonizing CES-1 activity in the same tissue.

DISCUSSION

In this study we report the first example of transcriptional activation by a Snail-type TF *in vivo*, as well as the first example of target sharing between Snail-type and bHLH TFs, as opposed to the widely reported Snail-type occlusion of bHLH DNA binding. Our data suggest that there are at least two types of CES-1 target genes (Figure 4c). First, there are the classical target genes that, within one cell or tissue, are either repressed by CES-1 or activated by bHLH-type TFs. In this case, these TFs compete for binding to their DNA recognition sequence. Second, there are targets that are activated by CES-1 and by bHLH-type TFs, but within different cells or tissues. In this case, these TFs do not compete for their DNA recognition sequence in the same tissue. To date, only two *in vivo* targets of any *C. elegans* Snail-type TF have been identified: *egl-1* (15) and the promoter of *B0507.1* (this study). CES-1 represses the first target, at least in *ces-1(gf)* animals that overexpress CES-1, but activates the second. The regulation by CES-1 of both *egl-1* and *PB0507.1* is highly tissue-specific; *egl-1* is repressed in the NSM sister cells in *ces-1(gf)* animals and *PB0507.1* is activated by CES-1 in the rectal gland, pharyngeal-intestinal valve, distal tip cells and spermatheca in wild-type animals. Although CES-1 expression is tissue-restricted in larvae and adults, it is broadly expressed in embryos (38; data not shown), suggesting that the expression patterns may be established early in development. No phenotype is known for *ces-1(lf)* mutants. Likewise, genome-scale RNAi studies have not yet revealed a biological function for either *ces-1* or *B0507.1* (WormBase). Thus, it is not yet conceivable to understand the transcriptional activation of *PB0507.1* by CES-1 in a phenotypic, or biological, context. Future studies are required to shed light on the mechanism of tissue-specific transcription activation and repression by CES-1; for instance, it may interact with tissue-specific transcriptional cofactors that either activate or repress expression, or may function in a combinatorial manner with other TFs.

At least two additional TFs contribute to the complex tissue-specific expression pattern conferred by the 300 bp




Figure 4. HLH-6 activates *B0507.1* promoter activity in the anterior pharyngeal gland cells. **(a)** Image of GFP expression in *PB0507.1::GFP;hlh-6(tm299)* animals. Left—GFP expression; center—DIC image; right—merge. **(b)** Quantification of GFP expression in *PB0507.1::GFP;hlh-6(tm299)* animals. **(c)** There are at least two types of CES-1 target genes. The first (target 1) are targets such as *egl-1* that contain one or more E-boxes (or CES-boxes) that are bound either by CES-1 or by a bHLH protein in the same cell (indicated by the gray rectangle), leading to gene repression or activation, respectively. The second (target 2) are targets like *PB0507.1* that contain one or more E-boxes (or CES-boxes) that are bound by both CES-1 and bHLH TFs, but in different cells (indicated by the green oval and blue hexagon). In the latter case, both TFs contribute to establishing a complex, multi-tissue expression pattern.

promoter of *B0507.1*. First, HLH-6 activates *PB0507.1* in the anterior pharyngeal gland cells. Although HLH-6 itself is expressed in both the anterior and posterior pharyngeal gland cells, it only regulates *PB0507.1* in the anterior pharyngeal gland cells. However, upon loss of HLH-6, the activity of *PB0507.1* was only partially reduced in this tissue. Both of these observations may result from *hlh-6(tm299)* not being a complete loss-of-function, or null allele. Expression in the anterior pharyngeal gland cells is reduced both in *hlh-6* mutants and in *PB0507.1* variants that carry mutations in the CES-box. In addition, it has previously been shown that HLH-6 regulates pharyngeal gland expression through an E-box sequence (28,41). These observations indicate that HLH-6 most likely activates *PB0507.1* by binding to the CES-box. Second, at least one as yet unidentified TF activates *PB0507.1* in the posterior pharyngeal gland cells. This TF has to act through a *cis*-regulatory element that is distinct from the CES-box because none of the *PB0507.1* mutants that we tested exhibited changes in GFP expression in these cells. We have previously identified another pharyngeal TF, ZTF-2, binding to *PB0507.1* (32). However, mutations in the *ztf-2* gene do not affect *PB0507.1* activity *in vivo* (data not shown). Thus, the remaining TF(s) that activate(s) *PB0507.1* may either be one of the two additional factors we identified by Y1H assays (ETS-5 and LIN-48) (32). However, these TFs are expressed in different tissues (WormBase). More likely, it is a TF that we have not retrieved in Y1H assays, for instance because the clone was not available (23), or because the TF binds DNA as an obligate

heterodimer (detection of which is not yet feasible in Y1H assays).

In sum, CES-1 can activate transcription *in vivo* and share a DNA target with bHLH proteins, albeit in different tissues. Future studies will determine whether these molecular functions are a common feature of Snail-type TFs and whether these TFs exhibit flexible functions in other systems as well. In addition, it will be interesting to see if binding site sharing in different tissues is a common feature of metazoan TFs.

SUPPLEMENTARY DATA

Supplementary Data are available at NAR Online.

ACKNOWLEDGEMENTS

The authors would like to thank Job Dekker and members of the Walhout lab for discussions and critical reading of the manuscript, Shohei Mitani for the *hlh-6(tm299)* allele and the *C. elegans* Genetics Center (CGC, supported by the NIH National Center for Research Resources) for additional strains.

FUNDING

National Institutes of Health [R01DK068429, R21DK071713 and R01GM082971 to A.J.M.W., R01GM069950 to B.C.]. J.G. is a Scholar of the Alberta Heritage Foundation for Medical Research and a Tier 2 Canada Research Chair in Developmental Genetics.

Canadian Institutes of Health Research Training Grant in Genetics, Child Health and Development (to R.B.S.). Funding for open access charge: R01GM082971.

Conflict of interest statement. None declared.

REFERENCES

1. Boulay, J.L., Dennefeld, C. and Alberga, A. (1987) The *Drosophila* developmental gene *snail* encodes a protein with nucleic acid binding fingers. *Nature*, **330**, 395–398.
2. Metzstein, M.M. and Horvitz, H.R. (1999) The *C. elegans* cell death specification gene encodes a Snail family zinc finger protein. *Mol. Cell*, **4**, 309–319.
3. Nakakura, E.K., Watkins, D.N., Schuebel, K.E., Sriuranpong, V., Borges, M.W., Nelkin, B.D. and Ball, D.W. (2001) Mammalian *scrunch*: a neural-specific Snail family transcriptional repressor. *Proc. Natl Acad. Sci. USA*, **98**, 4010–4015.
4. Hemavathy, K., Guru, S.C., Harris, J., Chen, J.D. and Ip, Y.T. (2000) Human Slug is a repressor that localizes to sites of active transcription. *Mol. Cell Biol.*, **26**, 5087–5095.
5. Hemavathy, K., Ashraf, S.I. and Ip, Y.T. (2000) Snail/Slug family of repressors: slowly going into the fast lane of development and cancer. *Gene*, **257**, 1–12.
6. Nieto, M.A. (2002) The Snail superfamily of zinc-finger transcription factors. *Nat. Rev. Mol. Cell Biol.*, **3**, 155–166.
7. Peinado, H., Olmeda, D. and Cano, A. (2007) Snail, ZEB and bHLH factors in tumour progression: an alliance against the epithelial phenotype? *Nat. Rev. Cancer*, **7**, 415–428.
8. Kasai, Y., Nambu, J.R., Lieberman, P.M. and Crews, S.T. (1992) Dorsal-ventral patterning in *Drosophila*: DNA binding of snail protein to the single-minded gene. *Proc. Natl Acad. Sci. USA*, **89**, 3414–3418.
9. Mauhin, V., Lutz, Y., Dennefeld, C. and Alberga, A. (1993) Definition of the DNA-binding site repertoire for the *Drosophila* transcription factor SNAIL. *Nucleic Acids Res.*, **21**, 3951–3957.
10. Fuse, N., Hirose, S. and Hayashi, S. (1994) Diploidy of *Drosophila* imaginal cells is maintained by a transcriptional repressor encoded by *escargot*. *Genes Dev.*, **8**, 2270–2281.
11. Bolos, V., Peinado, H., Perez-Moreno, M.A., Fraga, M.F., Esteller, M. and Cano, A. (2002) The transcription factor Slug represses E-cadherin expression and induces epithelial to mesenchymal transitions: a comparison with Snail and E47 repressors. *J. Cell Sci.*, **116**, 499–511.
12. Zeitlinger, J., Zinzen, R.P., Stark, A., Kellis, M., Zhang, H., Young, R.A. and Levine, M. (2007) Whole-genome ChIP-chip analysis of Dorsal, Twist, and Snail suggests integration of diverse patterning processes in the *Drosophila* embryo. *Genes Dev.*, **21**, 385–390.
13. Ip, Y.T., Park, R.E., Kosman, D., Bier, E. and Levine, M. (1992) The dorsal gradient morphogen regulates stripes of rhomboid expression in the presumptive neuroectoderm of the *Drosophila* embryo. *Genes Dev.*, **6**, 1728–1739.
14. Gray, S. and Levine, M. (1996) Short-range transcriptional repressors mediate both quenching and direct repression within complex loci in *Drosophila*. *Genes Dev.*, **10**, 700–710.
15. Thellmann, M., Hatzold, J. and Conradt, B. (2003) The Snail-like CES-1 protein of *C. elegans* can block the expression of the BH3-only cell-death activator gene *egl-1* by antagonizing the function of bHLH proteins. *Development*, **130**, 4057–4071.
16. Peinado, H., Ballestar, E., Esteller, M. and Cano, A. (2004) Snail mediates E-cadherin repression by recruitment of the Sin3A/histone deacetylase 1 (HDAC1)/HDAC2 complex. *Mol. Cell Biol.*, **24**, 306–319.
17. Nibu, Y., Zhang, H. and Levine, M. (1998) Interaction of short-range repressors with *Drosophila* CtBP in the embryo. *Science*, **280**, 101–104.
18. Qi, D., Bergman, M., Aihara, H., Nibu, Y. and Mannervik, M. (2008) *Drosophila* Ebi mediates Snail-dependent transcriptional repression through HDAC3-induced histone deacetylation. *EMBO J.*, **27**, 898–909.
19. Kataoka, H., Murayama, T., Yokode, M., Mori, S., Sano, H., Ozaki, H., Yokota, Y., Nishikawa, S.-I. and Kita, T. (2000) A novel Snail-related transcription factor Smuc regulates basic helix-loop-helix transcription factor activities via specific E-box motifs. *Nucleic Acids Res.*, **28**, 626–633.
20. Ellis, R.E. and Horvitz, H.R. (1991) Two *C. elegans* genes control the programmed deaths of specific cells in the pharynx. *Development*, **112**, 591–603.
21. Deplancke, B., Dupuy, D., Vidal, M. and Walhout, A.J.M. (2004) A gateway-compatible yeast one-hybrid system. *Genome Res.*, **14**, 2093–2101.
22. Deplancke, B., Vermeirssen, V., Arda, H.E., Martinez, N.J. and Walhout, A.J.M. (2006) Gateway-compatible yeast one-hybrid screens. *CSH Protocols*, doi:10.1101/pdb.prot4590.
23. Vermeirssen, V., Deplancke, B., Barrasa, M.I., Reece-Hoyes, J.S., Arda, H.E., Grove, C.A., Martinez, N.J., Sequerra, R., Doucette-Stamm, L., Brent, M. et al. (2007) Matrix and Steiner-triple-system smart pooling assays for high-performance transcription regulatory network mapping. *Nat. Methods*, **4**, 659–664.
24. Dupuy, D., Li, Q., Deplancke, B., Boxem, M., Hao, T., Lamesch, P., Sequerra, R., Bosak, S., Doucette-Stam, L., Hope, I.A. et al. (2004) A first version of the *Caenorhabditis elegans* Promoterome. *Genome Res.*, **14**, 2169–2175.
25. Walhout, A.J.M., Temple, G.F., Brasch, M.A., Hartley, J.L., Lorson, M.A., van den Heuvel, S. and Vidal, M. (2000) GATEWAY recombinational cloning: application to the cloning of large numbers of open reading frames or ORFeomes. *Methods Enzymol.: Chimeric Genes Proteins*, **328**, 575–592.
26. Praitis, V., Casey, E., Collar, D. and Austin, J. (2001) Creation of low-copy integrated transgenic lines in *Caenorhabditis elegans*. *Genetics*, **157**, 1217–1226.
27. Reece-Hoyes, J.S., Shingles, J., Dupuy, D., Grove, C.A., Walhout, A.J., Vidal, M. and Hope, I.A. (2007) Insight into transcription factor gene duplication from *Caenorhabditis elegans* Promoterome-driven expression patterns. *BMC Genomics*, **8**, 27.
28. Smit, R.B., Schnabel, R. and Gaudet, J. (2008) The HLH-6 transcription factor regulates *C. elegans* pharyngeal gland development and function. *PLoS Genet.*, **4**, e1000222.
29. Crooks, G.E., Hon, G., Chandonia, J.M. and Brenner, S.E. (2004) WebLogo: a sequence logo generator. *Genome Res.*, **14**, 1188–1190.
30. Ao, W., Gaudet, J., Kent, W.J., Muttumu, S. and Mango, S.E. (2004) Environmentally induced foregut remodeling by PHA-4/FoxA and DAF-12/NHR. *Science*, **305**, 1743–1746.
31. Walhout, A.J.M. (2006) Unraveling transcription regulatory networks by protein-DNA and protein-protein interaction mapping. *Genome Res.*, **16**, 1445–1454.
32. Deplancke, B., Mukhopadhyay, A., Ao, W., Elewa, A.M., Grove, C.A., Martinez, N.J., Sequerra, R., Doucette-Stam, L., Reece-Hoyes, J.S., Hope, I.A. et al. (2006) A gene-centered *C. elegans* protein-DNA interaction network. *Cell*, **125**, 1193–1205.
33. Vermeirssen, V., Barrasa, M.I., Hidalgo, C., Babon, J.A.B., Sequerra, R., Doucette-Stam, L., Barabasi, A.L. and Walhout, A.J.M. (2007) Transcription factor modularity in a gene-centered *C. elegans* core neuronal protein-DNA interaction network. *Genome Res.*, **17**, 1061–1071.
34. Martinez, N.J., Ow, M.C., Barrasa, M.I., Hammell, M., Sequerra, R., Doucette-Stamm, L., Roth, F.P., Ambros, V. and Walhout, A.J.M. (2008) A genome-scale microRNA regulatory network in *C. elegans* reveals composite feedback motifs that provide high information flow. *Genes Dev.*, **22**, 2535–2549.
35. Wolfe, S.A., Nekudova, L. and Pabo, C.O. (2000) DNA recognition by Cys2His2 zinc finger proteins. *Annu. Rev. Biophys. Biomol. Struct.*, **29**, 183–212.
36. Martinez, N.J., Ow, M.C., Reece-Hoyes, J., Ambros, V. and Walhout, A.J. (2008) Genome-scale spatiotemporal analysis of *Caenorhabditis elegans* microRNA promoter activity. *Genome Res.*, **18**, 2005–2015.
37. Hartley, J.L., Temple, G.F. and Brasch, M.A. (2000) DNA cloning using in vitro site-specific recombination. *Genome Res.*, **10**, 1788–1795.
38. Hatzold, J. and Conradt, B. (2008) Control of apoptosis by asymmetric cell division. *PLoS Biol.*, **6**, e84.

39. Ghai,V. and Gaudet,J. (2008) The CSL transcription factor LAG-1 directly represses *hlh-6* expression in *C. elegans*. *Dev. Biol.*, **322**, 334–344.
40. Krause,M., Park,M., Zhang,J.-M., Yuan,J., Harfe,B.D., Xu,S.-Q., Greenwald,I., Cole,M.D., Paterson,B.M. and Fire,A. (1997) A *C. elegans* E/Daughterless bHLH protein marks neuronal but not striated muscle development. *Development*, **124**, 2179–2189.
41. Raharjo,I. and Gaudet,J. (2006) Gland-specific expression of *C. elegans hll-6* requires the combinatorial action of three distinct promoter elements. *Dev. Biol.*, **302**, 295–308.