




Changes in Patient Reported Symptoms During the Natural Progression of Osteoarthritis

Matthew C. DeWolf MSIV, Patty Freedson PhD, David C. Ayers MD, Patricia D. Franklin MD MBA MPH

Department of Orthopedics and Physical Rehabilitation, University of Massachusetts Medical School



BACKGROUND

Arthritis is the leading cause of disability and second most costly chronic condition in the United States (1 & 2). Physical Activity is a challenge in patients with OA (3).

We quantitated the patient-reported changes in pain and function during the natural progression of OA and correlated these metrics with objective activity monitors.

MATERIALS & METHODS

50 patients who were undergoing non-operative management of OA were enrolled.

Visit Intervals: Baseline, 3 months, 6 months, 9 months

Data Collection:

- Basic Demographics
- Patient Reported Outcomes: SF-36, WOMAC, Charlson Co-Morbidity Index
- Objective Measures: ActiGraph and activPal*

* Worn for 7 days after visit

RESULTS

Basic Demographics

- ◆ Average Age: 57 years
- ◆ 80% had 1 or fewer medical co-morbidities
- ◆ 4% used an assistive device
- ◆ Average BMI: 33.85

Activity Monitor:

- Trend of increased % day sedentary, decreased steps/day. Table 1.

Patient - reported Measures:

- ◆ WOMAC Function: Average=68; NO change over time (0-100; moderate limitations)
- ◆ SF-36: Average PCS = 38; NO change over time (0-100; moderate limitations)
- ◆ If WOMAC Pain Score <80 (moderate pain); average SF36 PCS =36
- ◆ If WOMAC Pain Score >80 (mild pain); SF36 PCS =42.5

Table 1: Objective data from activPAL™ the initial 19 patients enrolled.

	%SED	%STAND	%STEP	STEPS	STEP RATE (steps per minute)	MVPA (minutes)
Baseline	62.8 ± 2.39	28.0 ± 1.98	9.2 ± 0.78	6446 ± 570	37.7 ± 0.94	52.6 ± 4.79
3-Mos	63.9 ± 2.40	28.0 ± 1.99	8.1 ± 0.79*	5359 ± 572*	36.6 ± 0.94*	42.6 ± 4.80*
6-Mos	62.6 ± 2.40	28.7 ± 1.98	8.7 ± 0.78	5868 ± 570	37.3 ± 0.94	47.1 ± 4.79*
9-Mos	67.1 ± 2.40*	24.7 ± 1.99	8.1 ± 0.79*	5322 ± 572*	37.3 ± 0.94	43.5 ± 4.80*


*Significantly different from baseline (p<0.05)

DISCUSSION

- ◆ Patient-reported function did not change over a 9-month period. However, preliminary objective activity data suggests a decline.
- ◆ Further analyses will correlate patient-reported measures with objective measures recorded by activity monitors to determine if objective monitors are preferable to detect early changes in activity due to OA.

Figure 1: Difference in SF-36 PCS between patients starting with a WOMAC Pain Score <80 and those starting with a WOMAC Pain Score >80, displayed at the 4 different study time points.

Correlation between WOMAC and SF-36 PCS



REFERENCES:

1. (CDC), Centers for Disease Control and Prevention. Prevalence of arthritis—United States, 1997. MMWR Morb Mortal Wkly Rep 2001. May 4; 50:334-6.

- Druss BG, Marcus SC, Olsson M, Pincus HA. The most expensive medical conditions in America. Health Affairs. 2002; 21:105-11.
- Centers for Disease Control and Prevention (CDC). Physical activity among adults with a disability—United States, 2005. MMWR Morb Mortal Wkly Rep 2007. Oct 5;56(39):1021-4.