

# eScholarship@UMassChan

## Does Functional Gain and Pain Relief After TKR and THR Differ by Patient Obese Status?

Item Type	Poster Abstract
Authors	Li, Wenjun;Lange, Jeffrey;Harrold, Leslie R;Allison, Jeroan J.;Lewis, Courtland;Bowen, Thomas;Ayers, David C.;Franklin, Patricia D
DOI	<a href="https://doi.org/10.13028/jzkr-jm42">10.13028/jzkr-jm42</a>
Rights	Copyright the Author(s)
Download date	2026-05-15 04:17:59
Item License	<a href="http://creativecommons.org/licenses/by-nc-sa/3.0/">http://creativecommons.org/licenses/by-nc-sa/3.0/</a>
Link to Item	<a href="https://hdl.handle.net/20.500.14038/27993">https://hdl.handle.net/20.500.14038/27993</a>

## Does Functional Gain and Pain Relief After TKR and THR Differ by Patient Obese Status?

Wenjun Li<sup>1</sup> PhD, Jeffrey Lange<sup>2</sup> MD, Leslie Harrold<sup>2</sup> MD MPH, Jeroan Allison<sup>3</sup> MD MS, Courtland Lewis<sup>4</sup> MD, Thomas Bowen<sup>5</sup> MD, David Ayers<sup>2</sup> MD, Patricia D. Franklin<sup>2</sup> MD MBA MPH

UMass Medical School <sup>1</sup>department of Preventive and Behavioral Medicine, <sup>2</sup>department of Orthopedics and Physical Rehabilitation, <sup>3</sup>department of Quantitative Health, Worcester, MA; <sup>4</sup>Hartford Hospital, Hartford CT, <sup>5</sup>Geisinger Medical Center, Danville, PA

Corresponding Author: Wenjun Li PhD, Associate Professor  
Department of Preventive and Behavioral Medicine, UMass Medical School  
Phone: 508-856-6574; Email: wenjun.li@umassmed.edu

**Introduction:** Obesity is an important predictor of functional status and pain after total knee (TKR) and total hip (THR) replacement. However, variations in pre-post TKR and THR changes in function and pain by obesity status remain to be examined.

**Material & Methods:** Pre- and 6 month post surgery data were collected on 2,964 primary TKR and 2,040 primary THR patients between 5/2011 and 3/2013. Data included demographics, comorbidities, operative joint pain severity based on the Knee Injury or Hip Disability and Osteoarthritis Outcome Score (KOOS/HOOS), WOMAC pain (higher is better), physical function (SF-36 PCS, higher is better), mental health (SF-36 MCS), and musculoskeletal burden of illness. Pre-post changes in PCS and pain were analyzed using descriptive statistics.

**Results:** TKR patients were average 67 years, 61% women, 93% whites, 13% under or normal weight, 33% overweight, 29% obese, 15% severely obese, 9% morbidly obese. Greater level of obesity was associated with lower PCS at baseline and 6 month, lower pain scores at baseline but larger improvement post-op. Pre to-6 month PCS did not differ by obesity status. At 6 months morbidly obese patients had slightly lower/worse pain score. THR patients were average 65 years, 62% women, 95% whites, 27% under/normal weight, 38% overweight, 23% obese, 9% severely obese, 4% morbidly obese. Greater level of obesity was associated with lower PCS at baseline and 6 month, poorer baseline pain score but larger improvement post-op. Mean changes in pre-to-6 month PCS did not differ by obesity status.

**Conclusion:** At 6 months after TKR, severely obese patients (BMI>35) reported improvements in both pain and function equal to or greater than patients with BMI<35. At 6 months after THR, all patients reported significant functional gains although patients with BMI>35 had lower mean functional gain than those with BMI<35. All patients reported excellent pain relief.