

eScholarship@UMassChan

Detection of CD8+ T cell Responses in Individuals with Long-term Type 1 Diabetes and Generation of Human CD8+ T Cell Lines Specific to Islet-associated Autoantigens

Item Type	Poster Abstract
Authors	Babon, Jenny Aurielle B;Harlan, David M.;Kent, Sally C
DOI	10.13028/sa9h-yy68
Rights	Copyright the Author(s)
Download date	2025-01-20 04:48:57
Item License	http://creativecommons.org/licenses/by-nc-sa/3.0/
Link to Item	https://hdl.handle.net/20.500.14038/27899

Detection of CD8+ T cell responses in individuals with long-term type 1 diabetes and generation of human CD8+ T cell lines specific to islet-associated autoantigens

Jenny Aurielle B. Babon, David M. Harlan and Sally C. Kent

Diabetes Center of Excellence, University of Massachusetts Medical School, Worcester, MA

Abstract:

Type 1 diabetes (T1D) is an autoimmune disease characterized by the activation of lymphocytes against insulin-producing β -cells in the pancreas. In humans, CD8+ T cells are predominantly found in sites of insulinitis and are considered to be one of the main drivers of β -cell destruction, thus indicating the need to analyze the frequency and function of these autoreactive CD8+ T cells. Peripheral blood mononuclear cells (PBMC) from individuals with long-term T1D were stained *ex vivo* for T cell surface markers and HLA-A2 pentamers containing known islet-associated epitopes to determine if there are autoreactive CD8+ T cells circulating in the periphery. All T1D donors tested had at least one detectable autoreactive CD8 T cell population and the frequencies of these autoantigen-specific T cells were comparable to previously published data from T1D individuals. We then developed a method of establishing CD8 T cell lines by co-culturing negatively isolated CD8 T cells and peptide-pulsed monocyte-derived dendritic cells from the PBMC of one T1D donor (A*02:01, A*33:01, B*14:02, B*40:01, DRB1*01:02, DRB1*04:04). We expanded a CD8 T cell line specific to the preproinsulin peptide PPI₁₅₋₂₄. This cell line produced IFN- γ and expressed CD107a in the presence of PPI₁₅₋₂₄-pulsed target cells, but not to an unrelated peptide or media alone. Using a similar approach, we were able to generate CD8 T cell lines from the same T1D donor that were cytotoxic to target cells pulsed with the autoantigens glutamic acid decarboxylase peptide (GAD65₁₁₄₋₁₂₃) and islet-specific glucose-6-phosphatase catalytic subunit-related protein peptide (IGRP₂₆₅₋₂₇₃). These autoreactive T cell lines can be utilized in *in vivo* assays using humanized mouse models to further understand the mechanism of β -cell destruction and disease progression. Studying the functionalities of these autoreactive T cells will also provide insights into identifying immune correlates to better assess both novel and existing immunotherapeutic strategies for T1D.

Contact info: Jenny Aurielle B. Babon
Jenny.Babon@umassmed.edu
508-856-1307